

# CONDITIONAL-MEAN HEDGING IN GAUSSIAN LONG-MEMORY MODELS WITH TRANSACTION COSTS

Tommi Sottinen

University of Vaasa

10th General AMaMeF Conference  
Padua, Italy, 25 June 2021

# ABSTRACT

We consider **INVERTIBLE GAUSSIAN VOLTERRA PROCESSES** and derive a formula for their **PREDICTION LAWS**. Examples of such processes include the fractional Brownian motions and the mixed fractional Brownian motions.

As an application, we consider **CONDITIONAL-MEAN HEDGING UNDER TRANSACTION COSTS** in Black-Scholes type pricing models where the Brownian motion is replaced with a more general invertible Gaussian Volterra process.

# OUTLINE

- 1 INVERTIBLE GAUSSIAN VOLTERRA PROCESSES AND THEIR PREDICTION
- 2 CONDITIONAL-MEAN HEDGING UNDER TRANSACTION COSTS FOR GEOMETRIC INVERTIBLE GAUSSIAN VOLTERRA MODELS
- 3 OPEN PROBLEMS

# OUTLINE

- 1 INVERTIBLE GAUSSIAN VOLTERRA PROCESSES AND THEIR PREDICTION
- 2 CONDITIONAL-MEAN HEDGING UNDER TRANSACTION COSTS FOR GEOMETRIC INVERTIBLE GAUSSIAN VOLTERRA MODELS
- 3 OPEN PROBLEMS

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

## DEFINITION (INVERTIBLE GAUSSIAN VOLTERRA PROCESS)

A centered continuous Gaussian process  $X$  on  $[0, T]$  with  $X_0 = 0$  is an **INVERTIBLE GAUSSIAN VOLTERRA PROCESS** if there exists a continuous Gaussian martingale  $M$  and kernels  $k$  and  $k^{-1}$  such that

$$\begin{aligned}X_t &= \int_0^t k(t, s) dM_s, \\M_t &= \int_0^t k^{-1}(t, s) dX_s.\end{aligned}$$

As an example, the **MIXED FRACTIONAL BROWNIAN MOTION**

$$X_t = aW_t + bB_t, \quad a, b \geq 0,$$

is an invertible Gaussian Volterra process.

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

Let  $\xi = (\xi', \xi'')$  be a jointly Gaussian **THING**. The **THEOREM OF GAUSSIAN CORRELATION** states that the **CONDITIONAL THING**  $\xi' | \xi''$  is a Gaussian **THING** with  $\xi''$ -measurable mean and deterministic covariance.

For **INVERTIBLE GAUSSIAN VOLTERRA PROCESSES** this theorem can be made practical. The key observation is:

$$\mathcal{F}_u^X = \mathcal{F}_u^M =: \mathcal{F}_u.$$

Let  $t \geq s \geq u$ . Denote

$$\begin{aligned}\hat{X}_t(u) &= E[X_t | \mathcal{F}_u], \\ \hat{r}(t, s | u) &= \text{Cov}[X_t, X_s, | \mathcal{F}_u].\end{aligned}$$

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

Now, for the conditional mean, since  $\mathcal{F}_u^X = \mathcal{F}_u^M$ , we have that

$$\hat{X}_t(u) = E[X_t | \mathcal{F}_u] = E \left[ \int_0^t k(t, s) dM_s \middle| \mathcal{F}_u^M \right].$$

Since Gaussian martingales have independent increments, we obtain

$$\begin{aligned} \hat{X}_t(u) &= E \left[ \int_0^u k(t, s) dM_s + \int_u^t k(t, s) dM_s \middle| \mathcal{F}_u^M \right] \\ &= \int_0^u k(t, s) dM_s. \end{aligned}$$

In order to write  $\hat{X}_t(u)$  as a linear transformation of the trajectory  $X_v$ ,  $v \in [0, u]$ , we note the following **TRANSFER PRINCIPLE**.

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

## THEOREM (TRANSFER PRINCIPLE)

$$\int_0^T g(t) dX_t = \int_0^T K_T^* g(t) dM_t,$$

where  $K_T^*$  is the operator defined by linearly extending the relation

$$K_T^* 1_{[0,t)}(s) = k(t, s).$$

If  $k$  is smooth enough and vanishes around the diagonal, then

$$K_T^* g(t) = \int_t^T g(s) k(ds, t).$$



# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

By the (inverse) transfer principle we can write

$$\begin{aligned}\hat{X}_t(u) &= \int_0^u k(t, s) dM_s \\ &= \int_0^u k(u, s) dM_s - \int_0^u [k(u, s) - k(t, s)] dM_s \\ &= X_u - \int_0^u (K_u^*)^{-1} [k(u, \cdot) - k(t, \cdot)](s) dX_s \\ &=: X_u - \int_0^u \Psi(t, s|u) dX_s.\end{aligned}$$

Note that if  $k^{-1}$  is smooth enough and vanishes around the diagonal, then

$$(K_T^*)^{-1} g(t) = \int_t^T g(s) k^{-1}(ds, t).$$

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

For the conditional covariance

$$\begin{aligned}\hat{r}(t, s|u) &= \text{Cov}[X_t, X_s | \mathcal{F}_u^X] \\ &= \text{E} \left[ \left( X_t - \hat{X}_t(u) \right) \left( X_s - \hat{X}_s(u) \right) \middle| \mathcal{F}_u^X \right]\end{aligned}$$

we have

$$\begin{aligned}X_t - \hat{X}_t(u) &= \int_0^t k(t, v) dM_v - \int_0^u k(t, v) M_v \\ &= \int_u^t k(t, v) dM_v, \\ X_s - \hat{X}_s(u) &= \int_0^s k(s, w) dM_w - \int_0^u k(s, w) M_w \\ &= \int_u^s k(s, w) dM_w.\end{aligned}$$

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

Consequently, since  $M$ , being a Gaussian martingale, has independent increments, we have that

$$\begin{aligned}\hat{r}(t, s|u) &= \mathbb{E} \left[ \int_u^t k(t, v) dM_v \int_u^s (s, w) dM_w \middle| \mathcal{F}_u^M \right] \\ &= \mathbb{E} \left[ \int_u^t k(t, v) dM_v \int_u^s (s, w) dM_w \right]\end{aligned}$$

Denote

$$q(t) = \text{Var}[M_t].$$

Note that  $q$  is also the **QUADRATIC VARIATION** of  $M$ .

Then, by the Itô isometry,

$$\hat{r}(t, s|u) = \int_u^{t \wedge s} k(t, v) k(s, v) dq(v).$$

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

Finally, we note that by the Itô isometry, we have

$$\begin{aligned}r(t, s) &:= \text{Cov}[X_t X_s] \\&= \mathbb{E} \left[ \int_0^t k(t, v) dM_v \int_0^s k(s, w) dM_w \right] \\&= \int_0^{t \wedge s} k(t, v) k(s, v) dq(v).\end{aligned}$$

Therefore,

$$\begin{aligned}\hat{r}(t, s|u) &= \int_u^{t \wedge s} k(t, v) k(s, v) dq(v) \\&= \int_0^{t \wedge s} k(t, v) k(s, v) dq(v) - \int_0^u k(t, v) k(s, v) dq(v) \\&= r(t, s) - \int_0^u k(t, v) k(s, v) dq(v).\end{aligned}$$

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

By the theorem of Gaussian correlation, we have the following:

## THEOREM (PREDICTION)

*The conditional process  $X_t|\mathcal{F}_u$ ,  $t \in [u, T]$  is Gaussian with mean and covariance given by*

$$\begin{aligned}\hat{X}_t(u) &= X_u - \int_0^u \Psi(t, s|u) dX_s, \\ \hat{r}(t, s|u) &= r(t, s) - \int_0^u k(t, v)k(s, v) dq(v).\end{aligned}$$

In what follows, we shall use the following short-hand

$$\hat{r}(t|u) := \hat{r}(t, t|u).$$

# INVERTIBLE GAUSSIAN VOLTERRA PROCESSES

Some final remarks on the invertible Gaussian Volterra processes:

- 1 If  $X$  is given by the covariance  $r$ , then the **VOLTERRA REPRESENTATION PROPERTY** means that there exist a measure  $q$  and a kernel  $k$  such that

$$r(t, s) = \int_0^{t \wedge s} k(t, v)k(s, v) dq(v).$$

- 2 The **INVERSE VOLTERRA REPRESENTATION PROPERTY** means that for all  $t \in [0, T]$  there exists a function  $g_t$  such that

$$K_T^* g_t = 1_{[0, t]}.$$

- 3 It is possible that  $X$  is continuous, but  $M$  is not.

# OUTLINE

- 1 INVERTIBLE GAUSSIAN VOLTERRA PROCESSES AND THEIR PREDICTION
- 2 CONDITIONAL-MEAN HEDGING UNDER TRANSACTION COSTS FOR GEOMETRIC INVERTIBLE GAUSSIAN VOLTERRA MODELS
- 3 OPEN PROBLEMS

# HEDGING UNDER TRANSACTION

We consider (imperfect) hedging under transaction costs in a discounted market model, where the the risky asset is given by

$$dS_t = S_t (\mu(t)dt + dX_t).$$

Let  $q$  be the quadratic variation of the fundamental martingale  $M$  associated with  $X$  via the kernel  $k$ . Then, the quadratic variation of  $X$  is (**UNDER SOME ASSUMPTIONS**)

$$\sigma^2(t) dt := k(t+, t) dq(t).$$

(We also assume that  $\sigma^2(t) \not\equiv 0$ , and note that that case is even simpler than the one studied here.)

We note that

$$\mathcal{F}_u := \mathcal{F}_u^S = \mathcal{F}_u^X = \mathcal{F}_u^M.$$



# HEDGING UNDER TRANSACTION

Let  $\pi = (\beta, \gamma)$  be a portfolio. Under continuous hedging without transaction costs its **SELF-FINANCING** value is

$$\begin{aligned}V_t &= \beta_t + \gamma_t S_t, \\V_t &= V_0 + \int_0^t \gamma_u dS_u.\end{aligned}$$

Suppose then that there are **PROPORTIONAL TRANSACTION COSTS**  $\kappa \in [0, 1)$  and (consequently) the trading takes place at times  $t_i$ ,  $i = 0, \dots, n$ . Let  $\pi^n = (\beta^n, \gamma^n)$  be a portfolio that is rebalanced at times  $t_i$ . Then the **SELF-FINANCING** value of  $\pi^n$  is

$$\begin{aligned}V_t^{n,\kappa} &= \beta_t^n + \gamma_t^n S_t, \\V_t^{n,\kappa} &= V_0^{n,\kappa} + \int_0^t \gamma_u^n dS_u - \kappa \int_0^t S_u d|\gamma_u^n|.\end{aligned}$$

# HEDGING UNDER TRANSACTION

Let the trading times  $t_i$  and the proportional transaction cost  $\kappa$  be fixed. Let  $f(S_T)$  be a European vanilla-type option. We are interested in solving the following imperfect hedging or tracking problem.

## DEFINITION (CONDITIONAL-MEAN HEDGING)

The strategy  $\pi^n$  is a **CONDITIONAL-MEAN HEDGE** of the option  $f(S_T)$  if for all  $t_i$

$$E[V_{t_{i+1}}^{n,\kappa} | \mathcal{F}_{t_i}] = E[V_{t_{i+1}} | \mathcal{F}_{t_i}],$$

where  $V$  is the value of the continuous-time perfect hedge without transaction costs.

# HEDGING UNDER TRANSACTION

Let  $\pi = (\beta, \gamma)$  be the replicating strategy of  $f(S_T)$  without transaction costs. Then, by the **FÖLLMER–ITÔ FORMULA**

$$\gamma_t = \frac{\partial g}{\partial x}(t, S_t),$$

where  $g(t, S_t) = V_t$  comes from the backward PDE

$$\begin{aligned} \frac{\partial g}{\partial t}(t, x) + \frac{1}{2}\sigma^2(t)x^2 \frac{\partial^2 g}{\partial x^2} &= 0, \\ g(T, x) &= f(x). \end{aligned}$$

Indeed, the derivation is the same as in the Black–Merton–Scholes case by using Föllmer integration, with the observation

$$(dS_t)^2 = S_t^2 \sigma^2(t) dt.$$

# HEDGING UNDER TRANSACTION

Let us then find out the conditional-mean values.

We denote

$$\begin{aligned}\hat{V}_{t_{i+1}}(t_i) &:= \mathbb{E} [V_{t_{i+1}} | \mathcal{F}_{t_i}], \\ \Delta \hat{V}_{t_{i+1}} &:= \hat{V}_{t_{i+1}}(t_i) - V_{t_i},\end{aligned}$$

$$\begin{aligned}\hat{V}_{t_{i+1}}^{n,\kappa}(t_i) &:= \mathbb{E} [V_{t_{i+1}}^{n,\kappa} | \mathcal{F}_{t_i}], \\ \Delta \hat{V}_{t_{i+1}}^{n,\kappa} &:= \hat{V}_{t_{i+1}}^{n,\kappa}(t_i) - V_{t_i}^{n,\kappa},\end{aligned}$$

$$\begin{aligned}\hat{S}_{t_{i+1}}(t_i) &:= \mathbb{E} [S_{t_{i+1}} | \mathcal{F}_{t_i}], \\ \Delta \hat{S}_{t_{i+1}} &:= \hat{S}_{t_{i+1}}(t_i) - S_{t_i},\end{aligned}$$

We will show that all these objects can be calculated explicitly by using the prediction formula for  $X_{t_{i+1}} | \mathcal{F}_{t_i}$ .

# HEDGING UNDER TRANSACTION

We note that if the initial portfolio  $\pi_0^n = (\beta_0^n, \gamma_0^n)$  is fixed, then  $V^{n,\kappa}$  can be recovered from the **CONDITIONAL-MEAN RECURSION**

$$\Delta \hat{V}_{t_{i+1}}^{n,\kappa}(t_i) = \Delta \hat{V}_{t_{i+1}}(t_i).$$

Let us then consider  $\hat{V}_{t_{i+1}}(t_i)$ :

$$\begin{aligned} \hat{V}_{t_{i+1}}(t_i) &= \mathbb{E} [V_{t_{i+1}} | \mathcal{F}_{t_i}] \\ &= \mathbb{E} [g(t_{i+1}, S_{t_{i+1}}) | \mathcal{F}_{t_i}] \\ &= \mathbb{E} \left[ g \left( t_{i+1}, S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2}\sigma^2(v)] dv + X_{t_{i+1}}} \right) \middle| \mathcal{F}_{t_i}^X \right] \\ &= \mathbb{E} \left[ g \left( t_{i+1}, S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2}\sigma^2(v)] dv + (X_{t_{i+1}} | \mathcal{F}_{t_i}^X)} \right) \right]. \end{aligned}$$

# HEDGING UNDER TRANSACTION

By using the **GAUSSIAN PREDICTION FORMULA**, we obtain

$$\begin{aligned}\hat{V}_{t_{i+1}}(t_i) &= \mathbb{E} \left[ g \left( t_{i+1}, S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2} \sigma^2(v)] dv + (X_{t_{i+1}} | \mathcal{F}_{t_i}^X)} \right) \right] \\ &= \int_{\mathbb{R}} g \left( t_{i+1}, S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2} \sigma^2(v)] dv + z} \right) \\ &\quad \times \varphi(z; \hat{X}_{t_{i+1}}(t_i), \hat{r}(t_{i+1} | t_i)) dz,\end{aligned}$$

where  $\varphi(\cdot; m, s^2)$  is the density of  $N(m, s^2)$  distribution.

Therefore, both  $\hat{V}_{t_{i+1}}(t_i)$  and

$$\begin{aligned}\Delta \hat{V}_{t_{i+1}}(t_i) &= \int_{\mathbb{R}} g \left( t_{i+1}, S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2} \sigma^2(v)] dv + z} \right) \\ &\quad \times \varphi(z; \hat{X}_{t_{i+1}}(t_i), \hat{r}(t_{i+1} | t_i)) dz - g(t_i, S_{t_i})\end{aligned}$$

are explicit.

# HEDGING UNDER TRANSACTION

Next, we note that

$$\begin{aligned}\hat{S}_{t_{i+1}}(t_i) &= \mathbb{E} [S_{t_{i+1}} | \mathcal{F}_{t_i}], \\ &= \int_{\mathbb{R}} S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2}\sigma^2(v)dv + z]} \varphi(z; \hat{X}_{t_{i+1}}, \hat{r}(t_{i+1}|t_i)) dz\end{aligned}$$

is explicit.

Consequently,

$$\begin{aligned}\Delta \hat{S}_{t_{i+1}}(t_i) &= \hat{S}_{t_{i+1}}(t_i) - S_{t_i} \\ &= \int_{\mathbb{R}} S_0 e^{\int_0^{t_{i+1}} [\mu(v) - \frac{1}{2}\sigma^2(v)dv + z]} \varphi(z; \hat{X}_{t_{i+1}}, \hat{r}(t_{i+1}|t_i)) dz - S_{t_i}\end{aligned}$$

is also explicit.

# HEDGING UNDER TRANSACTION

Finally, we note that

$$\begin{aligned}\hat{V}_{t_{i+1}}^{n,\kappa}(t_i) &= \mathbb{E} \left[ V_{t_{i+1}}^{n,\kappa} \middle| \mathcal{F}_{t_i} \right] \\ &= V_{t_i}^{n,\kappa} + \mathbb{E} \left[ \int_{t_i}^{t_{i+1}} \gamma_u^n dS_u - \kappa \int_{t_i}^{t_{i+1}} S_u |d\gamma_u^n| \middle| \mathcal{F}_{t_i} \right] \\ &= V_{t_i}^{n,\kappa} + \mathbb{E} \left[ \gamma_{t_i}^n (S_{t_{i+1}} - S_{t_i}) - \kappa S_{t_i} |\gamma_{t_i}^n| \middle| \mathcal{F}_{t_i} \right] \\ &= V_{t_i}^{n,\kappa} + \mathbb{E} \left[ \gamma_{t_i}^n S_{t_{i+1}} \middle| \mathcal{F}_{t_i} \right] - \mathbb{E} \left[ \gamma_{t_i}^n S_{t_i} \middle| \mathcal{F}_{t_i} \right] - \mathbb{E} \left[ \kappa S_{t_i} |\gamma_{t_i}^n| \middle| \mathcal{F}_{t_i} \right] \\ &= V_{t_i}^{n,\kappa} + \gamma_{t_i}^n \left( \hat{S}_{t_{i+1}}(t_i) - S_{t_i} \right) - \kappa S_{t_i} |\gamma_{t_i}^n|.\end{aligned}$$

Consequently,

$$\Delta \hat{V}_{t_{i+1}}^{n,\kappa}(t_i) = \gamma_{t_i}^n \Delta \hat{S}_{t_{i+1}}(t_i) - \kappa S_{t_i} |\gamma_{t_i}^n|.$$



# HEDGING UNDER TRANSACTION

We note that we have explicit (albeit horrible) formulas for all the quantities we need. Consequently, we have the following result:

## THEOREM (CONDITIONAL-MEAN HEDGING)

*Once the initial portfolio  $\pi_0^n = (\beta_0^n, \gamma_0^n)$  is fixed, the conditional-mean hedging portfolio for a European vanilla-type option  $f(S_T)$  can be calculated recursively from the explicit formulas for  $\Delta \hat{V}_{t_{i+1}}(t_i)$  and  $\Delta \hat{S}_{t_{i+1}}(t_i)$  by using the relations*

$$\begin{aligned}\Delta \hat{V}_{t_{i+1}}^{n,\kappa}(t_i) &= \Delta \hat{V}_{t_{i+1}}(t_i), \\ \Delta \hat{V}_{t_{i+1}}^{n,\kappa}(t_i) &= \gamma_{t_i}^n \Delta \hat{S}_{t_{i+1}}(t_i) - \kappa S_{t_i} |\gamma_{t_i}^n|.\end{aligned}$$

# OUTLINE

- 1 INVERTIBLE GAUSSIAN VOLTERRA PROCESSES AND THEIR PREDICTION
- 2 CONDITIONAL-MEAN HEDGING UNDER TRANSACTION COSTS FOR GEOMETRIC INVERTIBLE GAUSSIAN VOLTERRA MODELS
- 3 OPEN PROBLEMS

# OPEN PROBLEMS

- 1 Which Gaussian processes are invertible Gaussian Volterra processes with some fundamental martingale? **MY GUESS IS THAT ALL OF THE ARE.**
- 2 What is the natural initial condition for the conditional-mean hedge?  $V_0 = V_0^{n,\kappa}$  or  $V_0 = V_0^{n,\kappa} - \kappa S_0 |\gamma_0^n|$ ? Neither of these fixes the initial portfolio  $\pi_0^n = (\beta_0^n, \gamma_0^n)$ !
- 3 The conditional-mean hedging is natural from a tracking point of view. One can also consider the minimization problem

$$E [(V_T^{n,\kappa} - V_T)^2] \rightarrow \min!$$

This can be “solved” by using the **PREDICTION FORMULA**. The “solution” is a complete mess. It would be nice to know what is the connection to the conditional-mean hedge.

THANK YOU FOR LISTENING!

Any questions?

Any solutions to the open problems?